

Candida albicans Antigens

The diploid, unicellular yeast *Candida (C.) albicans* is an important commensal pathogen that colonizes the skin and mucosal surfaces of 30–70% of all humans (Dühring *et al.* 2015). In healthy individuals, *C. albicans* is controlled by the normal microbial flora, epithelial barriers, and the immune system. In situations when this homeostasis becomes imbalanced, e.g. burns leading to the destruction of epithelial barriers or a compromised immune system, a *C. albicans* infection, candidiasis, can develop (Dühring *et al.* 2015; Gow and Hube 2012). Superficial infections of the oral and vaginal mucosa are the most common forms of candidiasis. It is estimated that approximately 75% of all women suffer from vulvovaginal candidiasis at least once in their lifetime and up to 50% from recurrent infections. Especially in immunocompromised individuals, *C. albicans* can cause systemic infections by entering the blood stream (candidemia) and disseminating further through the body to infect virtually any organ (Gulati and Nobile 2016). Mortality rates of up to 50% have been reported for these systemic infections (Antinori *et al.* 2016; Dühring *et al.* 2015; Sudbery 2011).

Glucan 1,3-beta-glycosidase (Bgl2) is a cell wall localized 1,3-beta-glycosyltransferase, whose deletion negatively affects *C. albicans* virulence (Hartland *et al.* 1991; Sarthy *et al.* 1997). In 2006, Pitarch *et al.* analyzed sera from patients diagnosed with systemic candidiasis and found all to be positive for anti-Bgl2 antibodies.

C. albicans enolase is a glycolytic enzyme that catalyzes the conversion of 2-phosphoglycerate to phosphoenolpyruvate and is both, localized in the cytoplasm and cell wall-associated (Angiolella *et al.* 1996). Depending on the study, antibodies against this protein have been reported in approximately 50–90% of patients (Lain *et al.* 2007; Mitsutake *et al.* 1994; van Deventer *et al.* 1994).

The heat shock protein 70 (Hsp70) family comprises molecular chaperones involved in the synthesis and

transport of proteins. In *C. albicans*, the cytoplasmic Hsp70 or SSA1 also localizes to the cell wall and functions in the adhesion to host cells. Further, it reacts with patient sera in immunoassays (Bromuro *et al.* 1998; Chaffin *et al.* 1998; La Valle *et al.* 1995; López-Ribot *et al.* 1996).

C. albicans methionine synthase (Met6) is an essential, cobalamin-independent enzyme involved in the synthesis of methionine (Jones *et al.* 2004; Suliman *et al.* 2007) and has been reported as a major antigen in patients suffering from systemic candidiasis (Pardo *et al.* 2000; Pitarch *et al.* 2006).

DIARECT's *C. albicans* Bgl2, Enolase, Hsp70, and Met6 are produced in the baculovirus/insect cell expression system.




Figure: Immunodot analyses of sera from a blood donor (BD) and patients infected with *Candida albicans* (PS 1–4) for the presence of antibodies against recombinant *C. albicans* Bgl2, Enolase, Hsp70 and Met6. The proteins were printed on nitrocellulose membranes as indicated. Anti-human IgGMA (α -hu IgGMA) served as a positive control. Human serum albumin (HSA) and buffer (buffer ctrl) were used as negative controls.

References:

- Angiolella *et al.* (1996) J Infect Dis. 173 (3): 684-690
- Antinori *et al.* (2016) Eur J Intern Med. 34: 21-28
- Bromuro *et al.* (1998) Infect Immun. 66 (5): 2154-2162
- Chaffin *et al.* (1998) Microbiol Mol Biol Rev. 62 (1): 130-180
- Dühring *et al.* (2015) Front Microbiol. 6: 625
- Gow and Hube (2012) Curr Opin Microbiol. 15 (4): 406-412
- Gulati and Nobile (2016) Microbes Infect. 18 (5): 310-321
- Hartland *et al.* (1991) Proc Biol Sci. 246 (1316): 155-160
- La Valle *et al.* (1995) Infect immun. 63 (10): 4039-4045
- Lain *et al.* (2007) Clin Vaccine immunol. 14 (3): 318-319
- López-Ribot *et al.* (1996) Infect Immun. 64 (8): 3333-3340
- Mitsutake *et al.* (1994). J Clin Lab Anal. 8 (4): 207-210
- Pardo *et al.* (2000) Electrophoresis. 21 (13): 2651-2659
- Pitarch *et al.* (2006) Mol Cell Proteomics. 5 (1): 79-96
- Sarthy *et al.* (1997) Microbiology. 143 (Pt 2): 367-376
- Sudbery (2011) Nat Rev Microbiol. 9 (10): 737-748
- Suliman *et al.* (2007) Arch Biochem Biophys 467 (2): 218-226
- van Deventer *et al.* (1994) J Clin Microbiol. 32 (1): 17-23

In some countries the use of certain antigens in diagnostic tests may be protected by patents. DIARECT is not responsible for the determination of these issues and suggests clarification prior to use.

Ordering Information

44800	<i>Candida albicans</i> Bgl2	0.1 mg
44801		1.0 mg
45300	<i>Candida albicans</i> Enolase	0.1 mg
45301		1.0 mg
45400	<i>Candida albicans</i> Hsp70	0.1 mg
45401		1.0 mg
44900	<i>Candida albicans</i> Met6	0.1 mg
44901		1.0 mg

200325_Rev02

