

Tissue Transglutaminase (tTG; expressed in *E. coli*)

Antigen Specification

Product Number: 14400

Description:

Human tissue-type transglutaminase (tTG; Transglutaminase C, Transglutaminase H; EC 2.3.2.13). Recombinant antigen for in vitro research and manufacturing use only.

Immunological function:

Binds IgA- and IgG-type human auto-antibodies.

Origin:

Recombinant. Expressed in *E. coli* bacterial cells.

Expression construct:

cDNA coding for the long splice isoform of the human tissue-type transglutaminase fused to a hexa-histidine purification tag. By point mutation of the active center the catalytic transglutaminase activity has been eliminated, resulting in increased stability during storage and coating.

Biochemical tests:

SDS-PAGE (purity > 95%); Western blot with i: anti-tTG autoantibody positive sample; ii: monoclonal anti-His-tag antibody.

Calculated molecular weight:

78 kDa

Calculated isoelectric point:

pH 5.3

Immunological tests/Functionality:

Standard ELISA test (checkerboard analysis of negative and positive (anti-endomysium positive celiac disease sera) samples); immunodot analyses with positive/negative samples.

Recommended buffer/storage and handling conditions:

Recommendations for storage buffer: neutral to slightly alkaline pH and 20% glycerol as cryoprotective agent. Storage conditions: -70°C or below. Repeated freeze/thaw cycles should be avoided.

Coating concentration:

0.6-1.4 µg/mL (depending on the type of ELISA plate and coating buffer). Suitable for labeling of functional groups. Remark: very diverse coating protocols exist in the literature; DIARECT recommends extensive testing in assay development.

Copyright 2001 - 2020 DIARECT GmbH