

Product number:SZ04-0050 (50 mL)
SZ04-1000 (1000 mL)**Intended use:**

StabilZyme® conjugate/protein stabilizers protect the entire conjugate by preventing the loss of catalytic activity and maintaining the structural integrity of the protein in solution. These stabilizers allow for the storage of conjugated proteins at lower use concentrations, extended shelf life and increased signal-to-noise ratios for improved assay performance.

The StabilZyme NOBLE Stabilizer (BSA-Free) is optimized to maintain the conformation of antibody/antigen conjugates, antibody-coated particles and unmodified proteins that are commonly used in immunoassays. The key benefits of the BSA-free formulation are the elimination of protein interference and cross reactivity with the purpose of reducing erroneous results within immunoassay applications.

Adding the conjugate/protein directly into the NOBLE Stabilizer allows for stable storage of the conjugate at a working concentration and eliminates the need for subsequent dilutions.

Product stability, storage and specifications:

Product stability	Stable for 2.5 years from date of manufacture
Storage	Product should be stored at 2-8°C or at room temperature
Specification	Bovine Protein: None Product Buffer: MOPS pH: 6.0-7.0 Preservative: 0.02% methylisothiazolone and 0.02% bromonitrodioxane
Notes	Please note that SurModics Stabilization Products are shipped to customers at ambient temperature. Extensive stability studies have shown that prolonged storage at ambient temperature will not affect the product quality or efficacy.

Recommendations for use:

Aseptically pour off desired volumes of material needed for the application and allow the product to equilibrate at room temperature prior to use.

The following are general guidelines only.

- 1) Use StabilZyme NOBLE Stabilizer at 100% concentration for optimal stability and performance.
 - 10 mM MOPS or deionized water should be used if the product is diluted.
 - If a phosphate buffer is used, it is recommended the final concentration be no higher than a 5 mM buffer.
 - If necessary, adjust the pH of the solution to align with the properties of the protein for optimal performance.
- 2) Dilute the conjugate/protein/antibody coated particles to a working concentration in the StabilZyme NOBLE Stabilizer solution.
- 3) Use the diluted conjugate solution according to the lab defined protocol for the assay.
- 4) For optimal performance, store the conjugate solution at 2-8°C and protect from direct exposure to light.

Additional considerations:

An accelerated stability test is recommended to adequately assess the stability of the conjugate in-solution. Samples are stored at 37°C and 4°C for up to 90 days (longer if you prefer). At set intervals, the percent retained activity is calculated by dividing the 37°C sample optical density (O.D.) by the 4°C sample O.D. and plotted over time.

For technical assistance, email ivdtechsupport@surmodics.com

Related products:

In-Solution Protein Stabilizers & Diluents:
StabilZyme® HRP Conjugate Stabilizer (SZ02)
StabilZyme® SELECT Stabilizer (SZ03)
StabilZyme® Protein Free Stabilizer (SZPF)
Assay Diluent-HAMA Blocker (SM01)
Blockers/Stabilizers:
StabilGuard® Immunoassay Stabilizer-BSA-Free (SG01)
StabilCoat® Immunoassay Stabilizer (SC01)
Substrates:
BioFX® TMB One Component HRP Microwell Substrate (TMBW)
BioFX® TMB Supersensitive One Component HRP Microwell Substrate (TMBS)
BioFX® TMB Slow Kinetic One Component HRP Microwell Substrate (TMSK)
BioFX® TMB Enhanced HRP Membrane Substrate (ESPM)
BioFX® TMB One Component HRP Membrane Substrate (TMBM)
BioFX® ABTS One Component HRP Microwell Substrate (ABTS)

These products are for further manufacture and/or to be used as a component with in-vitro diagnostics immunoassays and are not intended for use in human or therapeutics purposes. Sales are without any seller's warranty or representation, expressed or implied, by usage or otherwise; no claims beyond replacement of unacceptable material or refund of purchase price shall be allowed. All claims must be made within 30 days following date of delivery.