

M2

Antigen Specification

Product Number: 18000

Description:

Human M2.
Mixture of E2/dihydrolipamide acyltransferase-subunits from 3 mitochondrial protein complexes: pyruvate dehydrogenase complex (PDC-E2); 2-oxo-glutarate dehydrogenase complex (OGDC-E2); branched chain 2-oxo-acid dehydrogenase complex (BCOADC-E2). Mixture contains equal mass of each protein component.
Recombinant antigen for in vitro research and manufacturing use only.

Immunological function:

Binds IgG-type human auto-antibodies.

Origin:

Recombinant. Expressed by recombinant baculovirus (*Autographa californica* multiple nuclear polyhedrosis virus; AcMNPV) infection of *Spodoptera frugiperda* Sf9 insect cells.

Expression constructs:

cDNAs coding for the mature forms of the human PDC-E2, OGDC-E2 and BCOADC-E2 proteins individually fused to a hexa-histidine purification tag.

Biochemical tests:

SDS-PAGE (purity > 75%); Western blot with i: anti-M2 autoantibody-positive sample; ii: monoclonal anti-His-tag antibody.

Calculated molecular weights:

PDC-E2	61 kDa
OGDC-E2	42 kDa
BCOADC-E2	47 kDa

Calculated isoelectric points:

PDC-E2	pH 5.8
OGDC-E2	pH 6.3
BCOADC-E2	pH 6.5

Immunological tests/Functionality:

Standard ELISA test (checkerboard analysis of positive/negative samples); immunodot analyses with positive/negative samples.

Recommended buffer/storage and handling conditions:

Recommendations for storage buffer: neutral to slightly alkaline pH. Storage conditions: -70°C or below. Repeated freeze/thaw cycles should be avoided.

Coating concentration:

0.4-0.8 µg/mL (depending on the type of ELISA plate and coating buffer). Suitable for labeling of functional groups.

Copyright 2006 - 2019 DIARECT AG